

## ORIGINAL ARTICLE

# Validation of self-reported energy intake by a self-administered diet history questionnaire using the doubly labeled water method in 140 Japanese adults

H Okubo<sup>1,2</sup>, S Sasaki<sup>2</sup>, HH Rafamantanantsoa<sup>3</sup>, K Ishikawa-Takata<sup>4</sup>, H Okazaki<sup>4</sup> and I Tabata<sup>4</sup>

<sup>1</sup>Department of Nutrition Science, Kagawa Nutrition University, Saitama, Japan; <sup>2</sup>Nutritional Epidemiology Program, National Institute of Health and Nutrition, Tokyo, Japan; <sup>3</sup>Department of Exercise and Sport Sciences, Shanghai Institute of Physical Education, Shanghai, PR China and <sup>4</sup>Health Promotion and Exercise Program, National Institute of Health and Nutrition, Tokyo, Japan

**Objective:** To validate reported energy intake (rEI) with a self-administered diet history questionnaire (DHQ) against total energy expenditure (TEE) by the doubly labeled water (DLW) method.

**Subjects:** A total of 140 healthy Japanese adults (67 men and 73 women) aged 20–59 years living in four areas in Japan.

**Methods:** Energy intake was assessed twice with DHQ over a 1-month period before and after TEE measurement (rEI<sub>DHQ1</sub> and rEI<sub>DHQ2</sub>, respectively). TEE was measured by DLW during 2 weeks (TEE<sub>DLW</sub>).

**Results:** Mean rEI<sub>DHQ1</sub> was lower than those of TEE<sub>DLW</sub> by  $1.9 \pm 2.4$  MJ/day (16.4%,  $P < 0.001$ ) for men and  $0.6 \pm 1.9$  MJ/day (6.0%,  $P < 0.01$ ) for women. In men and women together, 62 subjects (44%) were defined as underreporters (rEI<sub>DHQ1</sub>/TEE<sub>DLW</sub> < 0.84), 58 (41%) as acceptable reporters (0.84–1.16) and 20 (14%) as over-reporters (> 1.16). Pearson correlation coefficient was 0.34 for men and 0.22 for women. After adjustment for the dietary and non-dietary factors related to rEI<sub>DHQ1</sub>/TEE<sub>DLW</sub>, the correlation coefficient improved to 0.42 and 0.37, respectively.

**Conclusion:** The energy intake assessed with DHQ correlated low to modestly with TEE measured by DLW. In addition, DHQ underestimated energy intake at a group level. Caution is needed when energy intake was evaluated by DHQ at both individual and group levels.

*European Journal of Clinical Nutrition* (2008) **62**, 1343–1350; doi:10.1038/sj.ejcn.1602858; published online 1 August 2007

**Keywords:** doubly labeled water; energy intake; self-administered diet history questionnaire; validation; Japanese adults

## Introduction

Dietary intake estimates from self-administered dietary assessment methods such as questionnaires are commonly used in large-scale nutritional epidemiologic studies. Dietary assessment questionnaires have been developed for assessing habitual dietary intake and for ranking subjects according to

their dietary intake. However, they cannot entirely avoid reporting errors (Barrett-Connor, 1991), including not only random but also systematic errors (Black and Cole, 2001; Livingstone and Black, 2003), due to the fact that they are self-reported.

In validation studies, data from dietary assessment questionnaires have often been compared with data from reference methods such as weighed diet records or 24 h recall (Willett and Lenart, 1998). However, all these dietary assessment methods were based on self-reporting. Therefore, the errors of both the new and reference methods might be correlated each other. The doubly labeled water (DLW) method, which measures the total energy expenditure (TEE) of subjects in free-living situations, has made it possible to validate reported energy intake (rEI) with an external biomarker (Hill and Davies, 2001; Trabulsi and Schoeller, 2001). The error of the DLW method is independent of self-rEI error (Livingstone and Black, 2003). However,

Correspondence: Dr S Sasaki, Nutritional Epidemiology Program, National Institute of Health and Nutrition, 1-23-1 Toyama, Shinjuku-ku, Tokyo 162-8636, Japan.

E-mail: stssasak@nih.go.jp

Guarantor: S Sasaki.

**Contributors:** HO conducted the field management, data collection, statistical analysis and wrote the paper. SS conducted the study design and edited the paper. HHR, KIT and HO conducted data collection and IRMS analyses. IT conducted the overall management. All authors participated in the discussion and interpretation of the results, and in drafting and editing the paper.

Received 29 June 2006; revised 14 June 2007; accepted 18 June 2007; published online 1 August 2007

relatively few validation studies of food frequency questionnaires against the DLW method have appeared (Sawaya *et al.*, 1996; Andersen *et al.*, 2003; Subar *et al.*, 2003). Furthermore, no such studies have been reported in non-Western countries.

The purpose of the present study was to examine the validity of energy intake assessed with a self-administered diet history questionnaire (DHQ) (Sasaki *et al.*, 1998) in comparison with TEE, as measured by the DLW method in a Japanese population.

## Subjects and methods

### Study population

This study was conducted in four districts of Japan from May to August 2003. We invited 40 healthy subjects (20 men and 20 women) aged 20–59 years from each of the four areas to participate, and distributed five subjects equally in each sex and age class of 20–29, 30–39, 40–49 and 50–59 years. Details of study recruitment and enrollment were described previously (Ishikawa-Takata *et al.*, 2007). All subjects providing written informed consent were finally considered eligible for the study. The total number of participants was 157 (78 men and 79 women).

### Procedures

The study protocol was approved by the Ethics Committee of the National Institute of Health and Nutrition in Japan. The participants completed three visits over the study period and all participants completed the study. After recruitment, the participants were mailed an introductory letter and two dietary questionnaires including a DHQ, four physical activity questionnaires, and a supplemental questionnaire on lifestyle variables, and asked to fill them out and mail them back before the first visit (visit 1).

At visit 1, the participants had their questionnaires reviewed, their body weight and height measured and provided a baseline urine sample. At visit 2, on the morning following visit 1, they received a dose of DLW after an overnight fast. At visit 3, 14 days after visit 2, the participants brought urine samples and had their body weight and height measured.

After visit 3, the participants were mailed two dietary questionnaires including the DHQ, four physical activity questionnaires, supplemental questionnaire on lifestyle variables and diary about lifestyle during the period of TEE measurement.

All the collected questionnaires were checked by trained dietitians in each local center and again then in the study center. When missing answers, errors or both were found, the subjects were requested to answer the questions again.

### Dietary assessment methods

**Self-administered DHQ.** The DHQ is a validated 16-page structured questionnaire, which assesses dietary habits in the preceding 1-month period (Sasaki *et al.*, 1998, 2000). Details of the questionnaire, methods of calculating nutrients and validity are given elsewhere (Sasaki *et al.*, 1998, 2000). Briefly, the DHQ consists of seven sections; (1) general dietary behavior, (2) major cooking methods, (3) consumption frequency and amount of six alcoholic beverages, (4) consumption frequency and semiquantitative portion size of 121 selected food and nonalcoholic beverage items, (5) dietary supplements, (6) consumption frequency and amount of 19 staple foods (rice, bread, noodles and other wheat foods) and miso soup (fermented soybean paste soup), and (7) open-ended items for foods consumed regularly (=once/week), which are not listed in the question. The food and beverage items and portion sizes in the DHQ were derived primarily from the data in the National Nutrition Survey of Japan (Sasaki *et al.*, 1998) and several recipe books for Japanese dishes. Measures of energy and dietary intakes for food and beverage items and dietary supplements with energy (148 food items in total) were calculated using an ad hoc computer algorithm for the DHQ, which was based on the Standard Tables of Food Composition in Japan (Science and Technology Agency, 2000). Information on dietary supplements, such as tablet, powder and liquid, which contained few energy and on data from the open-ended questionnaire items were not used in the calculation of dietary intake.

### Anthropometric measures

Anthropometric measures were obtained at visits 1 and 3 by a single-trained study member. Body weight and height were measured to the nearest 0.1 kg and 0.1 cm, respectively, in subjects wearing light clothing and no shoes. Body mass index (BMI) was calculated as body weight (kg) divided by the square of body height ( $m^2$ ).

### Measurement of TEE with the DLW method

At visit 2, after a baseline urine sample was obtained, a single dose of approximately 0.06 g/kg body weight of  $^2H_2O$  (99.8 atom%, Cambridge Isotope Laboratories, MA, USA) and 0.14 g/kg body weight of  $H_2^{18}O$  (10.0 atom%, Cambridge Isotope Laboratories, MA, USA) was orally given to each subject via a drinking straw. After the dose administration, the subjects refrained from eating and drinking over a 4-h equilibration period (4 h sampling) for measurement of total body water. The second voided urine in the morning of day 1 (the day after the DLW dose) and day 14 (at the same time as the voiding on day 1) were collected for measurement of the isotopic ( $^2H$  and  $^{18}O$ ) elimination rate.

The procedure for specimen analysis and for subsequent data analyses was described previously (Ishikawa-Takata *et al.*, 2007). Briefly, the isotopic analyses were conducted

using the Isotope Ratio Mass Spectrometry (IRMS) *DELTA Plus* equipment (Thermo Electron Corporation, Bremen, Germany) and calibrated using Vienna Standard Mean Ocean Water (V-SMOW), 302B, and the Greenland Ice Sheet Precipitation (GISP) standard provided by the International Atomic Energy Agency. Each measurement of samples and the corresponding references was performed in duplicate. The average s.d. through the analyses were 0.5‰ for <sup>2</sup>H and 0.03‰ for <sup>18</sup>O.

TEE (kcal/day) calculation was performed using a modified Weir's formula Weir, 1949 based on rCO<sub>2</sub> (mol/day) and food quotient (FQ):

$$TEE = 3.9 \times (rCO_2 / FQ) + 1.1 \times (rCO_2)$$

FQ was derived from the dietary assessment data (g/day) of DHQ using an equation of Black *et al.* (1986). The average value of all subjects (0.867) was used for all subjects to estimate TEE.

#### Assessment of other variables possibly related to the rEI

Lifestyle, behavioral and psychological variables possibly related to the rEI were obtained from the four-page questionnaire as follows: educational attainment, alcohol drinking, history of diet experiences, desire for body weight change, and difference between ideal and measured body weight.

A physical activity level was calculated as TEE divided by basal metabolic rate (BMR). BMR was estimated according to the 6th Recommended Dietary Allowances for Japanese Ministry of Health Welfare (1999).

#### Statistical analysis

We excluded 17 subjects who was non-Japanese (*n* = 1), who was obese (*n* = 1), who did not complete at least first or second DHQ (*n* = 2), who had left more than 40 items blank in the questions regarding frequency for 121 selected food and beverage items in DHQ (*n* = 4), who rEI outside the range of 3.0–16.0 MJ/day (*n* = 2), or who did not provide sufficient urine sample volume (*n* = 7). Thus, 140 subjects (67 men and 73 women) were included in the present analysis.

As we monitored the body weight change during the assessment period of rEI by second DHQ (rEI<sub>DHQ2</sub>), we estimated EI (eEI) from TEE<sub>DLW</sub> with a correction for change in body energy store during the survey period (Bathalon *et al.*, 2000):

$$eEI = TEE + (\Delta wt \times 0.03)$$

where TEE is measured as MJ/day,  $\Delta wt$  is measured as g/day between visits 1 and 3, and 0.03 MJ/day (7 kcal/day) is the energy cost of weight change (Saltzman and Roberts, 1995). The eEI was used for the validation of rEI<sub>DHQ2</sub>. In contrast, this correction of change in body energy store was not considered for the validation of rEI<sub>DHQ1</sub> because of the lack of the monitoring.

The results were expressed as the mean and s.d. Mean differences between sexes and among methods were tested by the non-paired *t*-test and paired *t*-test, respectively. The Pearson and Spearman correlation coefficient was used to examine correlations between the test and the reference methods. Furthermore, the study participants were classified into tertiles of energy intake according to the distribution of

**Table 1** Characteristics of 140 Japanese men and women aged 20–59 years included in the analyses<sup>a</sup>

	Men ( <i>n</i> = 67)	Women ( <i>n</i> = 73)
Age (years)	39.4 ± 11.1	38.5 ± 10.4
Body height (cm)	169.3 ± 6.3	157.9 ± 6.1 <sup>e</sup>
Body weight (kg)	67.3 ± 9.7	53.9 ± 7.3 <sup>e</sup>
BMI (kg/m <sup>2</sup> ) <sup>b</sup>	23.3 ± 2.9	21.6 ± 2.7 <sup>e</sup>
< 18.5	5 (7)	10 (14) <sup>i</sup>
18.5–24.9	39 (58)	55 (75)
≥ 25.0	23 (34)	8 (11)
<i>Educational attainment</i>		
High school or less	28 (42)	23 (32) <sup>h</sup>
Technical or professional school	5 (7)	28 (38)
University or more	34 (51)	22 (30)
<i>History of diet experience<sup>c</sup></i>		
No	58 (87)	57 (78)
Yes	9 (13)	16 (22)
<i>Desire for weight change</i>		
Reduction	37 (55)	50 (68)
No change	20 (30)	20 (27)
Increase	10 (15)	3 (4)
Difference between ideal and measured body weight (kg) <sup>d</sup>	−4.2 ± 6.7	−4.5 ± 4.3
Frequency of alcohol intake (times/week)	2.6 ± 2.7	1.0 ± 1.9 <sup>e</sup>
Physical activity level	1.70 ± 0.21	1.69 ± 0.27
Body weight change during survey (g/day)	−23 ± 55 <sup>j</sup>	−2 ± 45 <sup>g</sup>
TEE <sub>DLW</sub> (MJ/day)	10.7 ± 1.7	8.3 ± 1.2 <sup>e</sup>
eEI <sub>DLW</sub> (MJ/day)	10.0 ± 2.1	8.2 ± 2.0 <sup>e</sup>
rEI <sub>DHQ1</sub> (MJ/day)	8.8 ± 2.4	7.7 ± 1.7 <sup>f</sup>
rEI <sub>DHQ2</sub> (MJ/day)	8.9 ± 2.5	7.4 ± 1.5 <sup>e</sup>

Abbreviations: BMI, body mass index; DHQ, diet history questionnaire; DHQ1, first measurement of DHQ before dose of DLW; DHQ2, second measurement of DHQ 2 weeks after dose of DLW; DLW, doubly labeled water method; eEI, estimated energy intake = TEE<sub>DLW</sub> + (body weight change during survey × 0.03); rEI<sub>DHQ</sub>, reported energy intake assessed with self-administered DHQ; TEE<sub>DLW</sub>, total energy expenditure measured by DLW.

<sup>a</sup>Mean ± s.d. or *n* (%).

<sup>b</sup>The categorization was based on the Japan Society for the Study of Obesity (Matsuzawa *et al.*, 2000).

<sup>c</sup>DiETING was defined as at least 2 kg intentional reduction of body weight within 1 month.

<sup>d</sup>Ideal body weight was evaluated by the following question: how many kilograms is your ideal body weight? Difference between ideal and measured body weight was calculated, as ideal body weight (kg)—measured body weight (kg), to evaluate the degree of desire for body weight change.

<sup>e–g</sup>Difference between sexes by non-paired *t*-test: <sup>e</sup>*P* < 0.001, <sup>f</sup>*P* < 0.01, <sup>g</sup>*P* < 0.05.

<sup>h</sup>Significant difference between sexes in all categories by  $\chi^2$  test: <sup>h</sup>*P* < 0.001, <sup>i</sup>*P* < 0.01

<sup>j</sup>Difference within sexes from 0 by paired *t*-test: *P* < 0.01.

the test and the reference methods, and the proportions of subjects classified into the same, adjacent or opposite tertiles were determined.

To evaluate the prevalence of under- or over-reporters, we calculated 95% confidence limits of  $rEI_{DHQ1}/TEE_{DLW}$  and  $rEI_{DHQ2}/eEI_{DLW}$  as a cutoff value proposed by Livingstone and Black (2003). Then, subjects with  $rEI_{DHQ1}/TEE_{DLW}$  and  $rEI_{DHQ2}/eEI_{DLW}$  smaller than 0.84 or larger than 1.16 were considered as under- or over-reporters, respectively.

A stepwise multiple regression analysis was performed to evaluate the influence of sociodemographic, lifestyle, behavioral and psychological factors on  $rEI_{DHQ1}/TEE_{DLW}$  and  $rEI_{DHQ2}/eEI_{DLW}$ , simultaneously. The following potential factors were entered into the model as the independent variables: age, BMI, body height, residential area, educational attainment, physical activity level, frequency of alcohol drinking, desire for body weight change, difference between ideal and measured body weight, and history of diet experience.

To examine the reproducibility, we compared mean rEIs between first and second DHQs (DHQ1 and DHQ2, respectively). Furthermore, the Pearson correlation coefficients were used to compare the rEIs assessed with DHQ1 and DHQ2.

All statistical analyses were performed using version 8.2 of the SAS software package (SAS Institute Inc., Cary, NC, USA). The test was considered significant at a *P*-value of <0.05.

## Results

Basic characteristics of the study subjects, the mean  $TEE_{DLW}$ , eEI, first and second measurements of rEI by the DHQ ( $rEI_{DHQ1}$  and  $rEI_{DHQ2}$ ) are shown in Table 1. Men had the higher BMI than women (23.3 versus 21.6 kg/m<sup>2</sup>, *P*<0.001).

Twenty-three of 67 men and eight of 73 women were overweight (BMI  $\geq 25$  kg/m<sup>2</sup>). This table also shows body weight change during the TEE measurement, between visits 1 and 3. Mean body weight in men, although not in women, significantly changed by  $-23 \pm 55$  g/day (*P*<0.01 by paired *t*-test). Mean  $rEI_{DHQ1}$  was significantly lower than mean  $TEE_{DLW}$  by  $1.9 \pm 2.4$  MJ/day (16.4%, *P*<0.001) for men and  $0.6 \pm 1.9$  MJ/day (6.0%, *P*<0.01) for women. Mean  $rEI_{DHQ2}$  was also significantly lower than mean  $eEI_{DLW}$  by  $1.1 \pm 2.7$  MJ/day (9.1%, *P*<0.001) for men and  $0.8 \pm 2.4$  MJ/day (4.6%, *P*<0.01) for women.

Table 2 shows reporting accuracy of energy intake assessed with DHQ expressed as  $rEI_{DHQ1}/TEE_{DLW}$  and  $rEI_{DHQ2}/eEI_{DLW}$ . The  $rEI_{DHQ1}/TEE_{DLW}$  and  $rEI_{DHQ2}/eEI_{DLW}$  was 0.84 and 0.91 for men and 0.94 and 0.95 for women, respectively, resulting in a significantly lower  $rEI_{DHQ1}/TEE_{DLW}$  ratio for men than for women (*P*<0.05). There was a wide range in reporting accuracy of DHQ1; 31 and 51% were identified as acceptable, and 58 and 32% as under-, and 10 and 18% as over-reporters for men and women, respectively.

The  $rEI_{DHQ1}$  and  $TEE_{DLW}$  were significantly correlated only for men (Pearson correlation coefficient = 0.34, Spearman correlation coefficient = 0.33), but not for women (0.22 and 0.16, respectively). Forty-one, 45 and 14% of the subjects were cross-classified into the same, the adjacent and the opposite tertiles of the respective distributions of  $rEI_{DHQ1}$  and  $TEE_{DLW}$ , respectively (Figure 1a). The results of the correlation between  $rEI_{DHQ2}$  and  $eEI_{DLW}$  were similar (Figure 1b).

Table 3 shows the results of multiple regression analysis with  $rEI_{DHQ1}/TEE_{DLW}$  and  $rEI_{DHQ2}/eEI_{DLW}$  as the dependent variables to examine the prediction of accuracy of reporting energy intake. For men, frequency of drinking alcohol, the difference between ideal and measured body weight, and history of diet experience correlated significantly and

**Table 2** Reporting accuracy of energy intake determined by the self-administered diet history questionnaire<sup>a</sup>

	DHQ1			DHQ2		
	All (n = 140)	Men (n = 67)	Women (n = 73)	All (n = 140)	Men (n = 67)	Women (n = 73)
Reporting accuracy <sup>b</sup>	0.89 ± 0.22	0.84 ± 0.21	0.94 ± 0.22 <sup>c</sup>	0.93 ± 0.30	0.91 ± 0.26	0.95 ± 0.33
Underreporters (n (%))	62 (44)	39 (58)	23 (32) <sup>d</sup>	64 (46)	30 (45)	34 (47)
Acceptable reporters (n (%))	58 (41)	21 (31)	37(51)	48 (34)	27 (40)	21 (29)
Overreporters (n (%))	20 (14)	7 (10)	13 (18)	28 (20)	10 (15)	18 (25)
Pearson's correlation coefficient	0.40 <sup>e</sup>	0.34 <sup>f</sup>	0.22	0.36 <sup>e</sup>	0.35 <sup>f</sup>	0.11
Spearman correlation coefficient	0.35 <sup>e</sup>	0.33 <sup>f</sup>	0.16	0.36 <sup>e</sup>	0.41 <sup>e</sup>	0.07

Abbreviations: DHQ1, first measurement of DHQ before dose of DLW; DHQ2, second measurement of DHQ 2 weeks after dose of DLW; DLW, doubly labeled water; eEI, estimated EI.


<sup>a</sup>Mean ± s.d. or n (%).

<sup>b</sup>Reporting accuracy was assessed as the ratio of energy intake to total energy expenditure ( $rEI_{DHQ1}/TEE_{DLW}$ ) and the ratio of energy intake to estimated energy intake ( $rEI_{DHQ2}/eEI_{DLW}$ ), respectively. eEI was determined by using a correction for change in body energy during the measurement period, as  $TEE_{\pm}(\text{body weight change during survey} \times 0.03)$ . Under-, acceptable, and over-reporters were defined as the ratio  $rEI_{DHQ}/TEE_{DLW}$  and  $rEI_{DHQ}/eEI_{DLW}$  <0.84, 0.84–1.16 and >1.16, respectively.

<sup>c</sup>Difference between sex by non-paired *t*-test: *P*<0.01.

<sup>d</sup>Significant difference between sexes in all categories by  $\chi^2$  test: *P*<0.01.

<sup>e,f</sup>Correlation coefficients between two methods: <sup>e</sup>*P*<0.001, <sup>f</sup>*P*<0.01.



**Figure 1** (a) Comparison of the first measurement of energy intake determined by the self-administered diet history questionnaire (rEI<sub>DHQ1</sub>) with total energy expenditure measured by the doubly labeled water method (TEE<sub>DLW</sub>) (■ = 67 men, ○ = 73 women). The dotted lines divide intake according to the tertiles of distribution. A straight line is  $y = x$ . Pearson and Spearman correlation coefficient was 0.40 and 0.35, respectively (both  $P < 0.001$ ). (b) Comparison of the second measurement of energy intake determined by the self-administered diet history questionnaire (rEI<sub>DHQ2</sub>) with estimated energy intake (eEI<sub>DLW</sub>) determined by a correction of body weight change during survey period, as  $TEE + (\Delta wt \times 0.03)$ , (■ = 67 men, ○ = 73 women). The dotted lines divide intake according to the tertiles of distribution. A straight line is  $y = x$ . Pearson and Spearman correlation coefficient was both 0.36 ( $P < 0.001$ ).

positively, and physical activity level negatively with rEI<sub>DHQ1</sub>/TEE<sub>DLW</sub>. For women, age and educational attainment correlated significantly and positively, and BMI negatively with rEI<sub>DHQ1</sub>/TEE<sub>DLW</sub>. We also conducted the same analysis with rEI<sub>DHQ2</sub>/eEI<sub>DLW</sub>. Body height, BMI and physical activity level significantly and negatively correlated with rEI<sub>DHQ2</sub>/eEI<sub>DLW</sub> for women. On the other hand, no factors attained the significance level for men.

The Pearson correlation coefficients between rEI<sub>DHQ1</sub> and TEE<sub>DLW</sub> slightly improved in both sexes after adjustment for

the above-mentioned related factors (0.42 for men and 0.37 for women).

We also examined reproducibility of energy intake between DHQ1 and DHQ2. The rEI<sub>DHQ2</sub> was significantly lower than rEI<sub>DHQ1</sub> for women (the difference was  $-0.3 \pm 1.1$  MJ/day,  $P = 0.03$ ), but not for men. The Pearson correlation coefficient between rEI<sub>DHQ1</sub> and rEI<sub>DHQ2</sub> was 0.79 for men and 0.76 for women.

## Discussion

To our knowledge, this is the first report in a non-Western country to validate energy intake estimated with a dietary assessment questionnaire against TEE measured by DLW method. Moreover, the sample size was relatively large compared to the previous studies with the same purpose and method (Sawaya *et al.*, 1996; Kroke *et al.*, 1999; Andersen *et al.*, 2003).

The mean rEI<sub>DHQ1</sub> was 11.0% less (16.4% for men and 6.0% for women) than the mean TEE<sub>DLW</sub>. Several validation studies have shown that dietary assessment instruments underestimated daily energy intake (Livingstone *et al.*, 1990; Hill and Davis, 2001). The degree of such error, under- or overestimation, has also been examined using TEE measured by the DLW method (Sawaya *et al.*, 1996; Kroke *et al.*, 1999; Andersen *et al.*, 2003; Livingstone and Black, 2003). Average underreporting in the previous studies between EI from dietary assessment questionnaires and TEE measured by DLW ranged from 10 to 38% (Sawaya *et al.*, 1996; Subar *et al.*, 2003), which depends on sample size and subjects (Trabulsi and Schoeller, 2001).

For the individual ranking, the rEI<sub>DHQ1</sub> significantly and positively correlated with TEE<sub>DLW</sub> ( $r = 0.40$ ,  $P < 0.001$ ), showing a correlation similar to or relatively higher than those observed in the previous studies ( $r = 0.06$ – $0.48$ ) (Kroke *et al.*, 1999; Bathalon *et al.*, 2000). Acceptable reporting was observed in 41% of the subjects, whereas 44% underreported and 14% over-reported. Underreporting of energy intake therefore seems to be a more serious problem than over-reporting.

In this study, the mean rEI<sub>DHQ1</sub>/TEE<sub>DLW</sub> ratio was significantly lower in men than in women. Further, the rate of underreporting was higher in men than in women. In a previous analysis of individual data from 21 studies, in contrast, the proportion of underreporters did not statistically differ between sexes (Black, 2000). In our previous study using semi-weighted diet records in 4 days  $\times$  4 seasons, the mean value of the ratio of rEI to BMR estimated from sex, age and body weight was not statistically different between sexes (Okubo *et al.*, 2006). In the DHQ, the portion sizes of food items are standardized regardless of sex, for example as 'one small cup'. The subjects then select the relative portion size from the five categories given except for rice, bread, noodles, other wheat foods and miso soup. This structure

**Table 3** Result of multiple regression analysis by stepwise procedure with the ratio of energy intake to total energy expenditure ( $rEI_{DHQ1}/TEE_{DLW}$  and  $rEI_{DHQ2}/eEI_{DLW}$ ) as dependent variables<sup>a</sup>

Independent variable <sup>b</sup>	Men (n = 67)						Women (n = 73)					
	DHQ1			DHQ2			DHQ1			DHQ2		
	Partial regression coefficient <sup>c</sup>	s.e. <sup>d</sup>	P-value	Partial regression coefficient <sup>c</sup>	s.e. <sup>d</sup>	P-value	Partial regression coefficient <sup>c</sup>	s.e. <sup>d</sup>	P-value	Partial regression coefficient <sup>c</sup>	s.e. <sup>d</sup>	P-value
Age (years)	—	—	—	—	—	—	0.005	0.002	0.04	—	—	—
BMI (kg/m <sup>2</sup> )	—	—	—	—	—	—	-0.036	0.009	<0.001	-0.049	0.015	<0.01
Body height (cm)	—	—	—	—	—	—	—	—	—	-0.016	0.006	0.02
Residential area	—	—	—	—	—	—	—	—	—	—	—	—
Educational attainment, (more than University versus high school or less as reference)	—	—	—	—	—	—	0.145	0.053	<0.01	—	—	—
Physical activity level	-0.356	0.120	<0.01	—	—	—	—	—	—	-0.480	0.154	<0.01
Frequency of drinking alcohol (times/week)	0.026	0.009	<0.01	—	—	—	—	—	—	—	—	—
Desire for body weight change	—	—	—	—	—	—	—	—	—	—	—	—
Difference between ideal and measured body weight (kg)	0.013	0.003	<0.01	—	—	—	—	—	—	—	—	—
History of diet experience (yes versus no as reference)	0.170	0.071	0.02	—	—	—	—	—	—	—	—	—

<sup>a</sup> $TEE_{DLW}$ , total energy expenditure measured by doubly labeled water method (DLW);  $rEI_{DHQ}$ , reported energy intake assessed with self-administered diet history questionnaire (DHQ); DHQ1, first measurement of DHQ before dose of DLW; DHQ2, second measurement of DHQ 2 weeks after dose of DLW. Reporting accuracy were assessed as the ratio of energy intake to total energy expenditure ( $rEI_{DHQ1}/TEE_{DLW}$ ) and the ratio of energy intake to estimated energy intake ( $rEI_{DHQ2}/eEI_{DLW}$ ), respectively. Estimated EI (eEI) was determined by using a correction for change in body energy during the measurement period, as  $TEE + (\text{body weight change during survey} \times 0.03)$ .

<sup>b</sup>See table 1 for the definition of each independent variable. Age (as a continuous variable), BMI (as a continuous variable), body height (as a continuous variable), residential area (Hokuriku, Shikoku, North Kyushu, and South Kyushu), educational attainment (high school or less, technical or professional school, or university or more), physical activity level (as a continuous variable), frequency of alcohol drinking (as a continuous variable), desire for body weight change (reduction, no change or increase), difference between ideal and measured body weight (as a continuous variable), history of diet experience (yes or no).

<sup>c</sup>Partial regression coefficient; change in dependent variable related to a 1-U change in independent variable.

<sup>d</sup>Standard error (s.e.) of the regression coefficient.

might have led to relative over- and underreporting of energy in women and men, respectively.

The  $rEI_{DHQ1}/TEE_{DLW}$  was significantly and independently correlated with several anthropometric and behavioral factors (Table 3). Several previous studies have already examined non-dietary factors, such as physiological (Zhang *et al.*, 2000; Livingstone and Black, 2003) and psychological (Johansson *et al.*, 1998; Bathalon *et al.*, 2000; Tooze *et al.*, 2004) factors associated with reporting accuracy of energy intake. After adjusting for these variables, the validity slightly improved (Pearson correlation coefficient was 0.42 for men and 0.37 for women). Therefore, these non-dietary factors are needed to consider when evaluating rEI.

This study has several limitations. First, FQ was derived from dietary assessment data by DHQ. Therefore, TEE was not theoretically independent of EI. Second, the surveyed period for the first measurement of EI by DHQ (DHQ1) was ahead of, and not overlapping with, TEE measurement by the DLW method. Third, we used the TEE as gold standard for the validation of DHQ1 without any consideration for a possible body weight change during the assessment period because of lack of the data. Fourth, we used the TEE with a correction for change in body weight during the survey period as gold standard for the validation of DHQ2, because the body weight has significantly changed in men. Fifth, the change in body composition, such as change in fat mass and fat-free mass, is probably the better indicator than the change in body weight for the correction of energy content for the study purpose. Sixth, the  $rEI_{DHQ1}$  was significantly lower than the  $rEI_{DHQ2}$  for women. Intentional or non-intentional intervention effect might have influenced dietary behaviors between the first and the second measurement. As shown in Table 3, the factors affecting reporting accuracy of energy intake were different between the two measurements. This may be one of the reasons. Seventh, we applied a two-point rather than multipoint method for the measurement of  $TEE_{DLW}$ . Eighth, the subjects were not randomly sampled from the general Japanese population. Moreover, the survey areas were not equally distributed over the country but were rather selected mostly from the Western parts of Japan.

In summary, the energy intake assessed with DHQ correlated low to modestly with TEE measured by DLW. In addition, DHQ underestimated energy intake at a group level. Caution is needed when energy intake was evaluated by DHQ at both individual and group levels.

## References

Andersen LF, Tomten H, Haggarty P, Lovo A, Hustvedt BE (2003). Validation of energy intake estimated from a food frequency questionnaire: a doubly labelled water study. *Eur J Clin Nutr* **57**, 279–284.

Barrett-Connor E (1991). Nutrition epidemiology: how do we know what they ate? *Am J Clin Nutr* **54** (Suppl 1), 182S–187S.

Bathalon GP, Tucker KL, Hays NP, Vincken AG, Greenberg AS, McCrory MA *et al* (2000). Psychological measures of eating behavior and the accuracy of 3 common dietary assessment methods in healthy postmenopausal women. *Am J Clin Nutr* **71**, 739–745.

Black AE (2000). The sensitivity and specificity of the Goldberg cut-off for EI:BMR for identifying diet reports of poor validity. *Eur J Clin Nutr* **54**, 395–404.

Black AE, Cole TJ (2001). Biased over- or under-reporting is characteristic of individuals whether over time or by different assessment methods. *J Am Diet Assoc* **101**, 70–80.

Black AE, Prentice AM, Coward WA (1986). Use of food quotients to predict respiratory quotients for the doubly labelled water method of measuring energy expenditure. *Hum Nutr Clin Nutr* **40C**, 381–391.

Hill RJ, Davies PS (2001). The validity of self-reported energy intake as determined using the doubly labelled water technique. *Br J Nutr* **85**, 415–430.

Ishikawa-Takata K, Tabata I, Sasaki S, Rafamantanantsoa HH, Okazaki H, Okubo H *et al.* (2007). Physical activity level in healthy free-living Japanese estimated by doubly labelled water method and International Physical Activity Questionnaire. *Eur J Clin Nutr* advance online publication, 23 May 2007; doi:10.1038/sj.ejcn.1602805.

Johansson L, Solvoll K, Bjørneboe G-EA (1998). Under- and over-reporting of energy intake related to weight status and lifestyle in a nationwide sample. *Am J Clin Nutr* **68**, 266–274.

Kroke A, Klipstein-Grobusch K, Voss S, Moseneder J, Thielecke F, Noack R *et al.* (1999). Validation of a self-administered food-frequency questionnaire administered in the European Prospective Investigation into Cancer and Nutrition (EPIC) Study: comparison of energy, protein, and macronutrient intakes estimated with the doubly labeled water, urinary nitrogen, and repeated 24-h dietary recall methods. *Am J Clin Nutr* **70**, 439–447.

Livingstone MB, Black AE (2003). Markers of the validity of reported energy intake. *J Nutr* **133** (Suppl 3), S895–S920.

Livingstone MB, Prentice AM, Strain JJ, Coward WA, Black AE, Barker ME *et al.* (1990). Accuracy of weighed dietary records in studies of diet and health. *BMJ* **300**, 708–712.

Matsuzawa Y, Inoue S, Ikeda Y, Sakata T, Saito Y, Sato Y *et al.* (2000). The judgment criteria for new overweight, and the diagnostic standard for obesity. *Obes Res* **6**, 18–28. (in Japanese).

Ministry of Health Welfare (1999). *Recommended dietary allowance for Japanese: dietary reference intakes*. 6th revised edn Ministry of Health and Welfare: Tokyo. (in Japanese).

Okubo H, Sasaki S, Hirota H, Notsu A, Todoriki H, Miura A *et al.* (2006). The influence of age and body mass index to relative accuracy of energy intake among Japanese adults. *Public Health Nutr* **9**, 651–657.

Saltzman E, Roberts SB (1995). The role of energy expenditure in energy regulation findings from a decade of research. *Nutr Rev* **53**, 209–220. Review.

Sasaki S, Ushio F, Amano K, Morihara M, Todoriki O, Uehara Y *et al.* (2000). Serum biomarker-based validation of a self-administered diet history questionnaire for Japanese subjects. *J Nutr Sci Vitaminol (Tokyo)* **46**, 285–296.

Sasaki S, Yanagibori R, Amano K (1998). Self-administered diet history questionnaire developed for health education: a relative validation of the test-version by comparison with 3-day diet record in women. *J Epidemiol* **8**, 203–215.

Sawaya AL, Tucker K, Tsay R, Willett W, Saltzman E, Dallal GE *et al.* (1996). Evaluation of four methods for determining energy intake in young and older women: comparison with doubly labeled water measurements of total energy expenditure. *Am J Clin Nutr* **63**, 491–499.

Science and Technology Agency (2000). *Standard Tables of Food Composition in Japan* 5th revised edn Printing Bureau, Ministry of Finance: Tokyo. (in Japanese).

Subar AF, Kipnis V, Troiano RP, Midthune D, Schoeller DA, Bingham S *et al.* (2003). Using intake biomarkers to evaluate the extent of

- dietary misreporting in a large sample of adults: the OPEN study. *Am J Epidemiol* **158**, 1–13.
- Tooze JA, Subar AF, Thompson FE, Troiano R, Schatzkin A, Kipnis V (2004). Psychosocial predictors of energy underreporting in a large doubly labeled water study. *Am J Clin Nutr* **79**, 795–804.
- Trabulsi J, Schoeller DA (2001). Evaluation of dietary assessment instruments against doubly labeled water, a biomarker of habitual energy intake. *Am J Physiol Endocrinol Metab* **281**, E891–899.
- Weir JBdV (1949). New methods for calculating metabolic rate with special reference to protein metabolism. *J Physiol* **109**, 1–9.
- Willett WC, Lenart E (1998). Reproducibility and validity of food-frequency questionnaires. In: Willett WC (ed). *Nutritional epidemiology*, 2nd edn. Oxford University Press: New York. pp 101–147.
- Zhang J, Temme EH, Sasaki S, Kesteloot H (2000). Under- and overreporting of energy intake using urinary cations as biomarkers: relation to body mass index. *Am J Epidemiol* **152**, 453–462.